



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 128 637**  
**A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 84302185.8

(22) Date of filing: 30.03.84

(51) Int. Cl.<sup>3</sup>: **C 12 P 13/22**  
// (C12P13/22, C12R1/13, 1/15)

(30) Priority: 13.04.83 JP 65010/83  
23.12.83 JP 243572/83

(43) Date of publication of application: 19.12.84  
Bulletin 84/51

(84) Designated Contracting States: DE FR GB IT

(71) Applicant: **AJINOMOTO CO., INC.**, 5-8,  
Kyobashi 1-chome, Chuo-ku, Tokyo 104 (JP)

(72) Inventor: **Shilo, Isamu**, No. 1-18-19, Sasuke,  
Kamakura-shi Kanagawa-ken (JP)  
Inventor: **Sugimoto, Shinichi**, No. 958, Kashimada,  
Saiwai-ku Kawasaki-shi Kanagawa-ken (JP)  
Inventor: **Kawamura, Kazuo**, No. 2-2-6, Nakase,  
Kawasaki-ku Kawasaki-shi Kanagawa-ken (JP)

(74) Representative: **Harrison, David Christopher et al**,  
**MEWBURN ELLIS & CO** 2/3 Cursitor Street, London  
EC4A 1BQ (GB)

(54) Process for the production of L-tryptophan by a fermentation process.

(57) A method for producing L-tryptophan by fermentation which comprises aerobically culturing in a culture medium a mutant of the genus *Corynebacterium* or genus *Brevibacterium* which is resistant to sulfaguanidine and capable of producing L-tryptophan, and recovering the L-tryptophan which has accumulated in the culture medium.

**EP 0 128 637 A1**

PROCESS FOR THE PRODUCTION OF L-TRYPTOPHAN  
BY A FERMENTATION PROCESS

The present invention relates to a process for the production of L-tryptophan (hereinafter referred to as tryptophan ) by a fermentation process.

Heretofore, there has been known, as a process for  
5 the production of tryptophan, a process which  
comprises producing tryptophan from a tryptophan  
precursor, that is, anthranilic acid, indole or 3-  
indolepyruvic acid. In contrast to this, the present  
inventors developed a process for the production of  
10 tryptophane by a direct fermentation process from a  
carbon source such as sugar by using a microorganism  
belonging to the genus Brevibacterium or the genus  
Corynebacterium and having resistance to tryptophan  
analogs such as 5-methyltryptophan, 5-fluoro-  
15 tryptophan etc. (Japanese Patent Publication No.  
18828/1973, French Patent Application Laid-open No.

2059715, Japanese Patent Application Laid-open No.  
162771/1980). It was also made clear that the amount  
of tryptophan accumulated is increased by imparting  
nutrient requirements for phenylalanine, tyrosine etc.  
5 and resistance to analogs of phenylalanine or tyrosine  
to these stains (Japanese Patent Publication No.  
18828/1973, French Patent Application Laid-open No.  
2059715) or by imparting thereto resistance to serine  
analogs (Japanese Patent Application Laid-open No.  
10 174096/1982).

The present inventors have studied for the purpose  
of developing a process for the production of  
tryptophane less expensively by these direct  
fermentation processes and, as a result, have  
15 discovered that by imparting resistance to  
sulfaguanidine, known as a sulfa drug, to the  
heretofore known tryptophan -producing microorganisms  
belonging to the genus *Corynebacterium* or the genus  
*Brevibacterium*, tryptophan is produced in much larger  
20 amounts than by the conventional tryptophan -producing  
microorganisms. The present invention has been  
accomplished as the result of our further study based  
on this discovery.

The microorganism used in the process for the production of tryptophan of the present invention is a mutant belonging to the genus *Corynebacterium* or the genus *Brevibacterium* and having the above-described  
5 properties necessary for the production of tryptophan, for example, having resistance to 5-methyltryptophan and having resistance to sulfaguanidine (it is a mutant having resistance to sulfaguanidine and capable of producing tryptophan.). By employing a strain having  
10 resistances to phenylalanine analogs, tyrosine analogs, azaserine, indolemycin, decoyinine etc. and nutrient requirements for L-phenylalanine, L-tyrosine, L-histidine, L-methionine etc. in addition to the resistance to sulfaguanidine, it is possible to  
15 increase the accumulation of tryptophan.

The parent strain for the mutant of the present invention is a microorganism belonging to the genus *Corynebacterium* or the genus *Brevibacterium* known as the so-called L-glutamic acid-producing micro-  
20 organism. Examples thereof include *Corynebacterium glutamicum* ATCC 13032, *Corynebacterium acetoacidophilum* ATCC 13870, *Corynebacterium lilium* ATCC 15990, *Brevibacterium flavum* ATCC 14067, *Brevibacterium divaricatum* ATCC 14020, *Brevibacterium lactofermentum*  
25 ATCC 13869, *Brevibacterium roseum* ATCC 13825.

The mutant employed in the present invention may be obtained by using the strain mentioned above as a parent strain, and subjecting it to mutating operations to impart thereto properties necessary for producing  
5 tryptophan, for example, resistance to 5-methyltryptophan and resistance to sulfaguanidine. In this case, the order of the operations for imparting these two properties is not restricted. The mutating operations may be effected in the conventional manner,  
10 for example, by irradiation with ultraviolet light or treatment with chemicals such as N-methyl-N'-nitro-N-nitrosoguanidine (hereinafter referred to as NG), nitrous acid.

Choosing *Corynebacterium glutamicum* AJ 12118 (FERM-  
15 P 7374, FERM-BP 478) and *Brevibacterium flavum* AJ 12022 (FERM-P 7034, FERM-BP 475) among the strains used in the present invention, their specific inducing methods and experimental examples showing the degrees of their resistance to sulfaguanidine are described below.

20 The mutants identified above by FERM-P numbers were originally deposited on December 19, 1983 and April 6, 1983 at the Fermentation Research Institute, Agency of Industrial Sciences and Technology, Ministry of International Trade and Industry (FRI), 1-3, Migashi  
25 1-Chome, Yatebe-machi, Tsukuba-gun, Ibaragi-ken 305, Japan, and were accorded the FERM-P number indicated

above. The mutant deposits were then converted into deposits under the Budapest Treaty on February 2, 1984, and were accorded the corresponding FERM-BP numbers.

Corynebacterium glutamicum FERM-P 1674 was used as  
5 a parent strain. The present strain is a strain which has been exemplified in Japanese Patent Publication No. 19037/1976 (Japanese Patent Application Laid-open No. 71194/1974) and, as its properties, described as having phenylalanine and tyrosine requirements and resistance  
10 to such phenylalanine and tyrosine analogs as p-aminophenylalanine, p-chlorophenylalanine and tyrosine hydroxamate, but further possesses tryptophan analog resistance. For example, when the degree of resistance to 5-methyltryptophan which is one of the tryptophan  
15 analogs to the FERM-P 1674 strain, was compared with a natural strain of Corynebacterium glutamicum ATCC 13032 using the method described in French Patent Application Laid-open No. 2059715, by inoculating test media containing 600 and 900  $\mu\text{g/ml}$  of 5-methyl-DL-  
20 tryptophan, respectively, with  $10^6$  cells and incubating at  $30^\circ\text{C}$  for 4 days, 1000 or more colonies were produced in either case with the FERM-P 1674 strain whereas there were only 122 and 0 colonies produced, respectively, with the natural strain.  
25 Thereafter, a sulfaguanidine-resistant strain was induced from the FERM-P 1674 strain. After treating

with 400  $\mu\text{g/ml}$  of NG at 30°C for 15 minutes (survival rate 12%), a plate medium was prepared by adding sulfaguanidine to the synthetic medium shown in Table 1 to a concentration where the parent strain cannot grow, i.e. 1000  $\mu\text{g/ml}$ .

Table 1

## Composition of the Synthetic Medium

Component		Concentration	
	Glucose	5	g/l
10	Urea	1.5	g/l
	Ammonium sulfate	1.5	g/l
	Potassium dihydrogenphosphate	3	g/l
	Potassium monohydrogenphosphate	1	g/l
	Magnesium sulfate	0.1	g/l
15	Calcium chloride	0.001	g/l
	Vitamin B <sub>1</sub> hydrochloride	100	$\mu\text{g/l}$
	d-Biotin	30	$\mu\text{g/l}$
	Minor metal elements*	1	ml/l
	L-Phenylalanine	100	mg/l
20	L-Tyrosine	100	mg/l
	Agar	20	g/l (pH 7.2)

\*The minor metal element solution used contains the following per liter:

8800 mg of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 970 mg of  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ , 270 mg of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ , 72 mg of  $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ , 88 mg of  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$  and 37 mg of  $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ .

After standing at 30°C for 10 days, the strains growing as colonies, i.e. the sulfaguanidine-resistant mutants, were harvested; of those, a mutant AJ 12118 (FERM-BP 478) excellently capable of producing

5    tryptophan (having sulfaguanidine resistance, tryptophan analog resistance, p-aminophenylalanine resistance, p-fluorophenylalanine resistance, tyrosine hydroxamate resistance, phenylalanine requirements and tyrosine requirements). This mutant produced

10   tryptophan in an amount 2.16 times that of the parent strain as shown in Example 1.

Thereafter, the results of the examination of the degree of resistance of this AJ 12118 strain to sulfaguanidine are shown in Table 2.

15       Sulfaguanidine was dissolved in the minimum medium shown in Table 1 at the concentrations shown in Table 2 to prepare plate media, which were then inoculated with about  $10^7$  cells of *Corynebacterium glutamicum* FERM-P 1674 and *Corynebacterium glutamicum* AJ 12118 (FERM-BP

20   478), respectively, grown in a complete medium (containing 10 g/l of yeast extract, 10 g/l of polypeptone, 5 g/l of sodium chloride, 5 g/l of glucose, 200 mg/l of L-methionine and 200 mg/l of L-tyrosine and at pH 7.0), incubated at 30°C for 3 days,

25   and the number of colonies produced was examined. The results are shown in Table 2.



Table 2

Degrees of Resistance of Strains to Sulfaguanidine

Concentration of Sulfa- guanidine ( $\mu\text{g/ml}$ )	Corynebacterium glutamicum		Brevibacterium flavum	
	FERM-P 1674	AJ 12118 (FERM-BP 478)	FERM-P 5907	AJ 12022 (FERM-BP 475)
0	+	+	+	+
1000	-	+		
1200			-	+

Note) In the table, (+) means that the number of colonies is 1000 or more, and (-) means that the number of colonies is zero.

Secondly, using *Brevibacterium flavum* AJ 11667  
 5 (FERM-P 5907) (having 5-fluorotryptophan resistance, p-fluorophenylalanine resistance, azaserine resistance, L-tyrosine requirements and L-methionine requirements) (Japanese Patent Application Laid-open No. 174096/1982) as a parent strain, this was treated with 200  $\mu\text{g/ml}$   
 10 of NG at 30°C for 15 minutes (survival rate 11%). Then, a plate medium was prepared by adding sulfaguanidine to the synthetic medium shown in Table 3 to a concentration where the parent strain cannot grow, i.e. 1200  $\mu\text{g/ml}$ , the mutation treated AJ 11667 was  
 15 applied thereto and, after standing at 30°C for 9 days,

strains growing as colonies, i.e. sulfaguanidine-resistant strains were harvested; of those, a mutant AJ 12022 (FERM-BP 475) excellently capable of producing tryptophan (having sulfaguanidine resistance, 5-fluorotryptophan resistance, p-fluorophenylalanine resistance, azaserine resistance, L-tyrosine requirements and L-methionine requirements) was harvested. This mutant produced tryptophan in an amount 68% higher than the parent strain as shown in

10 Example 2.

Table 3  
Composition of the Synthetic Medium

Component		Concentration	
	Glucose	20	g/l
	Ammonium sulfate	10	g/l
5	Potassium dihydrogenphosphate	1	g/l
	Magnesium sulfate	0.4	g/l
	Ferrous sulfate	10	mg/l
	Manganese sulfate	8	mg/l
	Sodium chloride	0.5	g/l
10	d-Biotin	50	µg/l
	Vitamin B <sub>1</sub> -HCl	200	µg/l
	L-Methionine	150	mg/l
	L-Tyrosine	100	mg/l
	L-Glutamic acid	30	mg/l
15	L-Threonine	100	mg/l
	Urea	3	g/l
	Agar	20	g/l
	(pH 7.2)		

Thereafter, the results of the examination of the  
20 degree of resistance of this AJ 12022 strain to  
sulfaguanidine are shown in Table 2.


Sulfaguanidine was dissolved in the synthetic  
medium shown in Table 3 at the concentrations shown in

Table 2 to prepare plate media, which were then inoculated with about  $10^7$  cells of *Brevibacterium flavum* AJ 11667 and *Brevibacterium flavum* AJ 12022, respectively, grown in a complete medium (containing 10 g/l of yeast extract, 10 g/l of polypeptone, 5 g/l of sodium chloride, 5 g/l of glucose, 200 mg/l of L-methionine and 200 mg/l of L-tyrosine and at pH 7.0), incubated at 30°C for 4 days, and the number of colonies produced was examined. The results are shown in Table 2.

As shown in Table 2, the mutants exhibited resistance to sulfaguanidine in contrast to the parent strains.

As can be seen from the experiments above, when resistance to sulfaguanidine is given to tryptophan producing microorganisms of the genus *Corynebacterium* or *Brevibacterium* by mutation, the productivity of tryptophane in the mutants obtained is remarkably increased. Mutants having resistance to sulfaguanidine also show resistance to other sulfa drugs than sulfaguanidine at the same time. Accordingly, mutants having resistance to other sulfa drugs other than sulfaguanidine are referred to herein as sulfaguanidine resistant mutants.

In this specification, the sulfa drugs mean those sulfa drugs which have the following general features and are generally known as antagonists to p-aminobenzoic acid. They contain in their molecule a group of the general formula:



and have anti-microbial action as a general rule. The anti-microbial action against usual wild strains is suppressed by addition of p-aminobenzoic acid.

Such sulfa drugs having the foregoing features include sulfapyridine, sulfathiazole, phthalylsulfathiazole, sulfadiazine, sulfaguanidine, sulfamethazine, sulfamerazine, sulfamethoxine, sulfamethomidine, sulfamethoxyridazine, sulfisomidine, sulfaoxazole, acetosulfamine, sulfanylamide, sulfisomezole, sulfaphenazole, sulfamethizole, sulfaethidole, sulfarazine, irgafen and irgamide.

The culture medium for producing tryptophane is not particularly restricted as is a conventional medium containing a carbon source, a nitrogen source, inorganic salts, and, if necessary, organic minor nutrients. As the carbon source, carbohydrates (glucose, fructose, or hydrolysates of starch, cellulose etc., molasses etc.), organic acids (acetic

acid, citric acid etc.), alcohols (glycerin, ethanol etc.) or hydrocarbons (normal paraffins etc.) may be employed. As the nitrogen source, ammonium sulfate, urea, ammonium nitrate, ammonium phosphate, ammonium  
5 chloride, ammonia gas etc. are employed, and as the inorganic salts, phosphates, magnesium salts, calcium salts, iron salts, manganese salts, and other minor metal salts are employed, according to the necessity. As the organic minor nutrients, if there are nutrient  
10 requirements, then appropriate amounts of the pertinent amino acids, vitamins, fatty acids, organic basic substances etc. are added, and, further, according to the necessity, amino acids, vitamins, AJIEKI (registered trademark; soy bean hydrolysate), yeast  
15 extract, peptone, Casamino acid, NZ amine, corn steep liquor etc. may be employed as growth promoting substances.

The incubating conditions may be conventional; for example, incubation may be conducted at pH 5-9 at a  
20 temperature of 20-40°C under aerobic conditions for 24-72 hours. If the pH is reduced during incubation, calcium carbonate, previously separately sterilized, is added, or neutralization is effected with alkali such as ammonia water, ammonia gas etc. On the other hand,  
25 where an organic acid is used as the carbon source, the increase in the pH is neutralized with a mineral acid or the organic acid.

Isolation and harvesting of tryptophan may be conducted in the conventional manner. It was confirmed with the obtained product that the Rf value on a paper chromatogram, the tryptophan specific reaction using  
5 Erlich reagent and the biological activity value by a microbiological quantitative method are in agreement with those of the tryptophan authentic product, and thus this was identified as tryptophan.

The quantitative assay of tryptophane was  
10 conducted according to the microbiological quantitative method using *Leuconostoc mesenteroides* (ATCC 8042).

The following examples are given for further understanding.

15 Example 1

A medium for producing tryptophan having the composition shown in Table 4 was allotted to 500 ml flasks, 20 ml per each, the microorganisms shown in Table 5 were inoculated in amounts of 1/3 slant  
20 respectively, and shake culture was conducted at 30°C for 72 hours. The amount of tryptophan produced in each culture broth was as shown in Table 5.

Table 4

Composition of the Medium for Producing Tryptophan		
Component	Concentration	
Molasses	100 (calculated as glucose)	g/l
5 Potassium dihydrogenphosphate	0.5	g/l
Potassium monohydrogenphosphate	0.5	g/l
Magnesium sulfate	0.25	g/l
Ammonium sulfate	20	g/l
10 Corn steep liquor	10	g/l
Calcium carbonate (separately sterilized)	20	g/l
L-Phenylalanine	200	mg/l
L-Tyrosine	175	mg/l
(pH 7.2)		

15

Table 5

Amount of Tryptophan Produced		
Strain	Sulfaguanidine Resistance	Amount of Tryptophane Produced (g/l)
20 FERM-P 1674	-	4.3
AJ 12118 (FERM-BP 478)	+	9.3

Note) In the table, (+) means that resistance is present, and (-) means that resistance is absent.



Example 2

A medium for producing tryptophan having the composition shown in Table 6 was allotted to 500 ml flasks, 20 ml per each, the microorganisms shown in Table 7 were inoculated in amounts of 1/3 slant, respectively, and shake culture was conducted at 30°C for 72 hours. The amount of tryptophan produced in each culture broth was as shown in Table 7.

Table 6

10	Composition of the Medium for Producing Tryptophan	
	Component	Concentration
	Glucose	130 g/l
	Ammonium sulfate	25 g/l
	Potassium dihydrogenphosphate	1 g/l
15	Fumaric acid	12 g/l
	Acetic acid	3 ml/l
	Manganese sulfate	8 mg/l
	d-Biotin	50 µg/l
	Vitamin B <sub>1</sub> ·HCl	2000 µg/l
20	L-Tyrosine	650 mg/l
	DL-Methionine	400 mg/l
	"AJIEKI"	50 ml/l
	Magnesium sulfate	1 g/l
	Calcium carbonate(Separately sterilized)	50 g/l
25	(pH 6.5)	

Table 7

Amount of Tryptophan Produced

Strain	Sulfaguanidine Resistance	Amount of Tryptophan Produced (g/l)
<i>Brevibacterium flavum</i>		
AJ 11667	-	9.1
AJ 12022 (FERM-BP 475)	+	15.3

Note) In the table, (+) means that resistance is present, and (-) means that resistance is absent.

CLAIMS:

1. A method for producing L-tryptophan by fermentation, which comprises: culturing aerobically in a culture medium a mutant of the genus *Corynebacterium* or genus *Brevibacterium* which is resistant to a sulfa drug  
5 and capable of producing L-tryptophan; and recovering the L-tryptophan which has accumulated in the culture medium.
2. A method according to Claim 1, wherein said mutant belongs to the species *Corynebacterium glutamicum*  
10 or *Brevibacterium flavum*.
3. A method according to Claim 2, wherein said mutant is *Corynebacterium glutamicum* FERM-BP 478 or *Brevibacterium flavum* FERM-BP 475 or their biological equivalents.
- 15 4. A method according to any one of preceding Claims, wherein the carbon source in the nutrient is sugar.



European Patent  
Office

# EUROPEAN SEARCH REPORT

0128637

Application number

EP 84 30 2185

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
A	FR-A-2 189 510 (KYOWA HAKKO KOGYO)  -----		C 12 P 13/22 // (C 12 P 13/22 C 12 R 1/13 C 12 R 1/15 )
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
			C 12 P C 12 D
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-10-1984	Examiner DESCAMPS J.A.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background U : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	